




Introduction to LBW for ALFALFA followup



Martha Haynes
UAT12 12.01.16

ALFALFA source codes



ALFALFA HI detections are coded according to:


Code 1	High quality sources, typically with $S/N > 6.5$
Code 2	Sources of lower S/N which are coincident with a probable OC of the same redshift (known from another source) => the "priors"
Code 3	Low S/N sources without identifiable OC
Code 4	Low S/N sources with a possible OC of unknown redshift
Code 5	Corresponding to Code 2, but of such low S/N or possible RFI contamination that they are untrustworthy
Code 6	Like OH megamasers at $0.16 < cz < 0.24$



Signal Extractor -- Introduction



- The signals are extracted by cross-correlations of a template with the spectra. => "matched-filter algorithm"
- More sensitive than peak-finding algorithms.
- Sensitive to total flux, not only peak flux
- Especially important for low mass systems
- Using FFT's, cross-correlations are computationally fast



Saintonge 2007 AJ 133, 2087

Slide: Amelie Saintonge




ALFALFA

Signal Extractor -- Application(2)



Saintonge 2007
AJ
133, 2087

The process is :



- Repeat for a range of widths of the template
 - e.g. 10 km/s – 600km/s
- Choose the width for which the convolution is maximised --> position of the signal
- Calculate the amplitude of the signal from the width



ALFALFA

Slide: Amelie Saintonge

High S/N



ALFALFA Catalog creator

X

File Imaging

(1) HI152947.8+260508_1532+25a.src
(2) HI152953.5+041105_1532+05c.src
(1) HI152956.8+072646_1532+07d.src
(1) HI153000.8+125922_1532+13b.src
(2) HI153003.0+261846_1532+27d.src
(4) HI153004.8+083711_1532+09c.src
(1) HI153009.3+252804_1532+25c.src
(4) HI153009.5+073415_1532+07d.src
(1) HI153011.6+055013_1532+05b.src
(5) HI153013.3+240023_1532+25d.src
(1) HI153020.8+074941_1532+07c.src
(3) HI153022.7+102807_1532+11b.src
(1) HI153023.9+271608_1532+27a.src
(1) HI153031.3+065630_1532+07b.src
(1) HI153031.9+144209_1532+15b.src
(1) HI153032.8+251550_1532+25b.src
(1) HI153034.0+050850_1532+05c.src
(1) HI153035.0+264447_1532+27b.src
(1) HI153037.2+272859_1532+27c.src

STATUS

- ◇ (0) No status
- ◇ (1) Detection
- ◇ (2) Prior
- ◇ (3) Marginal
- ◇ (4) Low StN
- ◇ (5) Prior-
- ◇ (9) HVC

Mark \ Unmark

HI152947.8+260508

V50,W50: 2019.5 68.2+/- 5.5 km/s
V20,W20: 2019.8 107.4+/- 5.5 km/s
Vcen: 2016.2+/- 2.7 km/s
V,W Gauss: 0.0 0.0+/- 0.0 km/s
Stot(profile, P): 2.02+/- 0.07 Jy km/s
Stot(profile, G): 0.00+/- 0.00 Jy km/s
Map Stot: 1.92+/- 0.00 Jy km/s
meanS, peakS: 11.7 27.6 mJy
S/N P: 23.1 12.4 11.7 29.0
S/N G: 0.0 0.0 0.0 0.0
Cont: 13. mJy
Status Code: 1

(l,b)= (40.49, 54.74) degrees
Cen_cell: 152949.3+260515 [2000]
Opt pos: 152948.2+260516 [2000]
dRA: -1.07685 sec
dDec: 1.33 arcsec
Ellipse: 4.0 x 3.5 PA= -18.
Isophote: 880. mJy km/s
Map Smax: 1759. mJy km/s
rms: 2.35 mJy

AGC727130

MODIFY PARAMETERS

Optical Coordinates

Signal/Noise

Modify SNR

cz Err Stat/Sys /

Width Err Stat/Sys /


AGC Number cz(opt)

Save Changes


SDSS data:

Acquire SDSS


View AGC Info



Select Isophote:



SDSS ObjID: 587739382067167453 specObjID: 609061544193425408
petroMag(Err): u= 17.91(0.09) g= 17.06(0.02) r= 16.82(0.01) i= 16.65(0.04) z= 16.89(0.11)
extinct: u= 0.27 g= 0.20 r= 0.14 i= 0.11 z= 0.08
petroR50_r: 6.11 petroR90_r: 13.46 expAB_r: 0.33



SDSS Navigator

SkyView

5

NED



ALFALFA

Low S/N



ALFALFA Catalog creator

File Imaging

- (1) HI153606.5+100300_1540+09c.src
- (2) HI153607.3+250823_1532+25c.src
- (2) HI153608.9+252713_1532+25c.src
- (1) HI153609.1+054730_1532+05c.src
- (1) HI153609.5+125056_1532+13d.src
- (3) HI153614.7+153955_1532+15a.src
- (1) HI153618.3+054010_1532+05c.src
- (3) HI153618.3+090430_1532+09b.src
- (2) HI153621.7+120758_1540+13d.src
- (5) HI153623.1+264000_1532+27c.src
- (2) HI153628.6+121235_1532+13d.src
- (4) HI153639.9+124212_1540+13c.src
- (1) HI153641.4+090122_1540+09c.src
- (1) HI153645.9+075008_1532+09b.src
- (2) HI153646.1+074430_1532+07d.src
- (2) HI153647.1+252419_1532+25c.src
- (1) HI153650.0+261027_1532+27c.src
- (1) HI153650.8+035153_1540+05c.src
- (1) HI153651.2+050639_1540+05c.src

STATUS

- ◇ (0) No status
- ◇ (1) Detection
- ◇ (2) Prior
- ◇ (3) Marginal
- ◇ (4) Low StN
- ◇ (5) Prior-
- ◇ (9) HVC

Mark \ Unmark

HI153606.5+100300

V50,W50: 10243.3 213.0+/- 9.0 km/s
 V20,W20: 10245.3 222.4+/- 9.0 km/s
 Vcen: 10248.6+/- 4.5 km/s
 V,W Gauss: 0.0 0.0+/- 0.0 km/s
 Stot(profile, P): 1.00+/- 0.08 Jy km/s
 Stot(profile, G): 0.00+/- 0.00 Jy km/s
 Map Stot: 0.70+/- 0.00 Jy km/s
 meanS, peakS: 4.0 5.8 mJy
 S/N P: 7.0 2.1 2.7 12.3
 S/N G: 0.0 0.0 0.0 0.0
 Cont: 15. mJy
 Status Code: 1

(l,b)= (17.20, 47.67) degrees
 Cen_ell: 153606.6+100252 [2000]
 Opt pos: 153606.8+100248 [2000]
 dRA: 0.18605 sec
 dDec: -3.79 arcsec
 Ellipse: 4.3 x 3.8 PA= 0.
 Isophote: 359. mJy km/s
 Map Smax: 719. mJy km/s
 rms: 2.20 mJy

AGC715089

MODIFY PARAMETERS

Optical Coordinates


Signal/Noise

cz Err Stat/Sys /


Width Err Stat/Sys /

AGC Number cz(opt)


SDSS data:



Select Isophote:



No SDSS information acquired for this file.



SDSS Navigator

SkyView

NED



ALFALFA

"Dark" galaxy candidate



ALFALFA Catalog creator

File Imaging

(1) HI152835.1+061437_1532+07a.src

STATUS

- (0) No status
- (1) Detection
- (2) Prior
- (3) Marginal
- (4) Low StN
- (5) Prior-
- (9) HVC

Mark \ Unmark

HI152835.1+061437

V50, W50: 509.1 24.3+/- 15.9 km/s
V20, W20: 500.2 50.6+/- 15.9 km/s
Vcent: 498.6+/- 8.0 km/s
V, W Gauss: 0.0 0.0+/- 0.0 km/s
Stot(profile, P): 0.33+/- 0.05 Jy km/s
Stot(profile, G): 0.00+/- 0.00 Jy km/s
Map Stot: 0.22+/- 0.00 Jy km/s
meanS, peakS: 3.8 9.9 mJy
S/N P: 6.3 5.5 4.3 7.0
S/N G: 0.0 0.0 0.0 0.0
Cont: 9. mJy
Status Code: 1

(l,b)= (10.99, 47.27) degrees
Cen_ell: 152834.9+061429 [2000]
Opt pos: 000000.0+000000 [2000]
dRA: ***** min
dDec: ***** arcmin
Ellipse: 3.9 x 2.8 PA= 20.
Isophote: 123. mJy km/s
Map Smax: 246. mJy km/s
rms: 2.29 mJy

AGC258472

tantalizing, no oc

MODIFY PARAMETERS




Optical Coordinates

Signal/Noise

cz Err Stat/Sys /

Width Err Stat/Sys /

AGC Number cz(opt)



No SDSS information acquired for this file.

SDSS Navigator

SkyView

NED



ALFALFA

Identifying Optical Counterparts




ALFALFA source centroids good to $\sim 18''$ (depends on S/N)

ALFALFA catalogs include:

- the HI centroid position
- the position of the most probable OC
- OC's SDSS PhotoObjID and SpecObjID (where applicable)


Of 15855 sources in $\alpha.40$:

- 1013 have no OC
- 844 of those could be HVCs (or LG minihalos)
- 199 ($< 2\%$) extragalactic
- Of those, < 50 are "isolated"



ALFALFA

ALFALFA advantage for finding the OC



Centroiding accuracy
goes roughly as
 $\text{HPFW}(\text{PSF})/(\text{S/N})$

Suppose HIPASS detects a
source at $\text{S/N} \sim 6$ near 3000 km/s
in this field. The position error
box will have a radius of $\sim 2.5'$.

The opt counterpart could be gal
#1, 2, 3, 4, 5 or 6.

**ALFALFA will detect the
same source with $\text{S/N} \sim 50$**

and the Arecibo beam is $\frac{1}{4}$ as
wide as the Parkes one

→ The same source will have
an ALFALFA position error
of $\sim 0.1'$



ALFALFA

Dark galaxies




- In agreement with previous results, ALFALFA finds that **fewer than 2%** of (clearly extragalactic) HI sources cannot be identified with an optical counterpart.
- The majority of objects without OC's are found near to galaxies with similar redshifts.
- There are few interesting cases to be confirmed (work in progress).



ALFALFA

"Dark" object in a group



HI peak with no optical/marginal UV: almost dark?

MH, RG (CU), Cannon (Macalester), Salzer (Indiana)



ALFALFA

HI114310.1+141330
J114310.3+141328.9

HI121850.1+123621
J121851.3+123549.9

HI122022.6+121136
J122022.9+121108.9

HI122710.8+155407
J122711.8+155349.9


HI123506.6+123100
J123507.99+123020

HI12440
J124409

HI122942.6+094202
J122942.96+094152

HI12301
J123019

HI125401.5+092700
J125402.09+092648.9




Ongoing deeper B,R band imaging of HI
detections with no OC

Salzer, Adams

OH Megamasers: OHMs

- Arise in interacting/merging galaxy systems.
 - When galaxies merge, gas clouds close to their nuclei are shocked and heated by the collision, and the emission from certain molecules especially OH is strongly amplified.
- Since this stimulated emission is like the more familiar laser but occurs in the **microwave** region of the electromagnetic spectrum, it is called a "**maser**".
- When galaxies collide, the emission is millions of times stronger than in normal galaxies, hence the term "**megamaser**".
- Such objects are also typically (ultra) luminous in the far-infrared.





Ultraluminous Infrared Galaxy Arp 220 HST • NICMOS


PRC97-17 • ST ScI OPO • June 9, 1997
R. Thompson (University of Arizona),
N. Scoville (California Institute of Technology) and NASA




18 cm OHMs



18 cm OH lines




- First molecular lines detected in radio.
- 18 cm lines formed by "lambda doubling" hyperfine structure
- 1612, 1665, 1667 and 1720 MHz



Pacholczyk, *Radio Astrophysics*




Redshifted 18 cm OHMs in ALFALFA



- $f_5 = 1665.4018$
 - $f_7 = 1667.3590$
- In OHMs, $S(f_7) > S(f_5)$

$$f_{\text{obs}} = f_{\text{rest}} / (1+z)$$

ALFALFA: 1340-1430 MHz,
corresponding, for OH, to
 $0.166 < z < 0.244$



- Emission at $f > 1422$ MHz (blueshifted if HI)
- Emission associated with OC in $0.1666 < cz < 0.244$
- Emission with no OC




OHM candidates in a.40



Table 4
OH Megamaser Candidates

AGC	OHM Coords (J2000) (hh mm ss.s+dd mm ss)	Opt. Coords (J2000) (hh mm ss.s+dd mm ss)	z_{opt}	z_{OH}	v_{21} (km s^{-1})	F_{OH} (Jy km s^{-1})	S/N	rms (mJy)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
102708	000337.0+253215	000336.1+253204		0.169	-1335	0.91	5.7	2.33
102850	002958.8+305739	002958.2+305832		0.172	-596	0.46	6.7	2.09
181310	082311.7+275157	082312.7+275138	0.16783	0.168	-1551	2.17	15.9	2.18
228040	124540.5+070337	124545.7+070347		0.172	-624	0.33	5.1	2.11



AGC 181310:

- Previously discovered OHM by Darling & Giovanelli (2001)
- Coincides with SDSS J083212.61+275139.8 at $z=0.167830$
- Also IRAS 08201+2801 and 5C 07.206

We need to confirm the others, and a few more low SNR sources.

AGC 181310: confirmed OHM



ALFALFA Catalog creator

File Imaging

(1) HI082310.4+275152_0820+27a.src

(4) HI082312.2+040114_0820+05c.src

(9) HI082316.6+280500_0820+27a.src

(1) HI082318.0+273035_0820+27b.src

(1) HI082323.9+244012_0820+25b.src

(1) HI082327.6+092410_0820+09c.src

(2) HI082327.8+251745_0820+25d.src

(1) HI082328.1+270816_0820+27b.src

(1) HI082329.1+150918_0820+15a.src

(1) HI082330.6+100300_0820+09b.src

(3) HI082333.3+150300_0820+15b.src

(1) HI082336.3+250100_0820+25b.src

(9) HI082337.4+280010_0820+27a.src

(1) HI082341.5+035346_0820+05a.src

(1) HI082350.9+095100_0820+09c.src

(1) HI082351.3+144525_0820+15a.src

(1) HI082352.7+280610_0820+27d.src

(1) HI082356.5+113334_0820+11c.src

(2) HI082358.2+040952_0820+05c.src

STATUS

◇ (0) No status

◇ (1) Detection

◇ (2) Prior

◇ (3) Marginal

◇ (4) Low StN

◇ (5) Prior-

◇ (9) HVC

Mark \ Unmark

HI082310.4+275152

V50,W50: -1515.7 193.5+/- 26.2 km/s

V20,W20: -1491.5 290.6+/- 26.2 km/s

Vcent: -1534.0+/- 13.1 km/s

V,W Gauss: 0.0 0.0+/- 0.0 km/s

Stot(profile, P): 2.17+/- 0.07 Jy km/s

Stot(profile, G): 0.00+/- 0.00 Jy km/s

Map Stot: 1.69+/- 0.00 Jy km/s

meanS, peakS: 9.2 18.9 mJy

S/N P: 15.9 5.1 8.6 29.0

S/N G: 0.0 0.0 0.0 0.0

Cont: 23. mJy

Status Code: 1

(l,b)= (195.30, 31.31) degrees

Centr_ell: 082311.7+275157 [2000]

Opt pos: 082312.7+275138 [2000]

dRA: 1.02306 sec

dDec: -19.48 arcsec

Ellipse: 4.3 x 3.1 PA= -75.

Isophote: 875. mJy km/s

Map Smax: 1749. mJy km/s

rms: 2.18 mJy

AGC181310

clear, undisputable OHM, w/oc of matching z

cz(HI) = -1518 km/s

MODIFY PARAMETERS

Optical Coordinates 082312.7+275138

Signal/Noise 15.9 Modify SNR


cz Err Stat/Sys 13.1000 / 0.000000

Width Err Stat/Sys 26.200068 / 0.000000


AGC Number 181310 cz(opt) 50215

Save Changes SDSS data: Checked no photo


View AGC Info



Select Isophote: 874



No SDSS information acquired for this file.



SDSS Navigator

SkyView

NED



ALFALFA sensitivity & completeness




Figure 12. Distribution of $\alpha.40$ extragalactic sources in the profile width vs. integrated flux density ($\log W_{50}$ – $\log S_{21}$) plane. The upper panel shows the distribution of Code 1 detections only, while the lower panel shows the same for the whole $\alpha.40$ catalog, including Code 1 (blue symbols) and Code 2 (green symbols) detections. In both panels, the solid red line corresponds to the 90% completeness limit, while the red dash-dotted line corresponds to the 50% (“sensitivity limit”) and the red dotted line to the 25% (“detection limit”) completeness limits. See Section 6 for the analytical expressions for the plotted limits, as well as for an explanation of the derivation method.

- We want to integrate longer on the low S/N sources.
- Even on high S/N sources, we want to verify they are real.
- Point at OC if there is one or the HI centroid if not
“Targeted observations”
- LBW has a single horn (“pixel”) but higher gain and lower T_{sys} than ALFA

Position-switched observing




The signals we are trying to take are billions of times weaker than the radio noise contributed by the receiver, electronics, antenna, cosmic microwave background and the sky overall.

Somehow, we have to subtract off all those unwanted contributions to find our signal.

We assume that a random position in the sky does not contain an HI line source at the exact same velocity as our target source.

We observe such a position, but track it over the exact same Az, ZA as our observations of the target source. => ON-OFF pair



Estimating how long we integrate



The radiometer equation for our observations

$$S_{\text{rms}} = \frac{(T_{\text{sys}}/G)}{\sqrt{2\Delta f_{\text{ch}} t_s f_t}},$$

For LBW,
 $T \sim 30\text{K}$
 $G \sim 11 \text{ K/Jy}$

Δf : is the bandwidth per channel

t_s : is the effective integration time, in secs

f_t : accounts for the degree of smoothing, the technique applied for bandpass subtraction, clipping losses, etc.

The factor of 2 under the square root comes from the fact that we average the two independent polarizations.

See Giovanelli + 2005, AJ 130, 2598



A2669 LBW followup proposal



Targeted LBW observations of selected ALFALFA sources:

1. "Dark" galaxy candidates: high quality (Code 1) detections with no OC and not associated with known tidal debris fields
2. OH megamaser (OHMs) candidates: either at large blueshift or coincident (within centroiding accuracy) of OC of appropriate cz.
3. "Low mass dwarf candidates": low signal-to-noise ratio sources at low cz (< 1000 km/s)
4. Statistical samples of low S/N signals possibly associated with optical galaxies

ALFALFA: effective integration time of 40 seconds/beam
LBW: 3 minutes ON-source

