



What HI tells us about Galaxies and Cosmology



Martha Haynes

UAT17 17.06.14



ALFALFA: A Census of Gas-bearing Galaxies



- A galaxy is a gravitationally bound object that consists of billions (and billions) of stars, gas clouds (of varying temperature and density = interstellar medium), dust clouds (mixed with the gas), and (so it seems), 90% dark matter.
- Optical surveys, like the Sloan Digital Sky Survey, detect the **stellar component** of galaxies.
- ALFALFA is designed to detect the **cool** (not hot; not cold) atomic gas in and near galaxies.
- ALFALFA is a **blind** survey; we observe the whole area of sky, whether or not we think/know there is an optical galaxy there.
- ALFALFA is a **spectroscopic** survey; not only do we detect the HI line flux, we also measure its frequency (velocity) and the width of the HI line (a measure of rotational velocity).



The HI 21 cm line @ 1.42 GHz

HI : Why do we care ?




- Easy to detect, simply physics → cold gas mass
- Good index of SF fertility → future SF
- Comparative HI content ⇒ HI deficiency
- Excellent tracer of host dynamics → dark matter
- Useful Cosmology tool → HI mass function, HI velocity width function, Tully-Fisher relation, "dark" galaxies(?)
- Interaction/tidal/merger tracer
- Can be **dominant baryon form** in low mass galaxies
- **ALFALFA**: A census of HI in the local universe



Hydrogen in the Interstellar Medium



- **HI** is the designation astronomers use for neutral hydrogen atoms in space.
- It is estimated that **4.4%** of the visible matter in our galaxy is HI. $\Rightarrow \sim 4.8 \times 10^9 M_{\odot}$.
- The **fraction** of interstellar space filled with HI clouds is 20% to 90%.



Full-Sky Map at 1420 MHz
Shows distribution of HI


Atomic vs molecular gas

- Estimates for molecular hydrogen, H_2 , vary - 1.2 to $3.5 \times 10^9 M_{\odot}$.
- H_2 tends to concentrate in a small number of giant gas clouds.
- Found principally in the inner region, where most of the SF also occurs.
- Stars form when **molecular clouds** collapse (usually)
- The **atomic** gas is the "fuel reservoir".

HI emission from galaxies



- Under most circumstances, the total H I mass can be derived from the integrated line profile; that is, the line **flux density** (integrated over all frequencies where there is signal) is **proportional** to the **number of hydrogen atoms**.
- The frequency (velocity) spread of the line reflects the **velocities of the gas** atoms, not quantum mechanics => hence the width of the line tells about the motions of the gas (rotation within the galaxy or turbulence, expansion, etc)



$$\begin{aligned} \int F dV &\rightarrow \text{HI mass} \\ V &\rightarrow \text{Distance} \\ \Delta V &\rightarrow \text{Mass} \end{aligned}$$

Rest frequency 1420.4058 MHz

Clues from the HI line


- **Redshifts** (\Rightarrow distances via Hubble's Law)
- **HI mass and distribution** (for extended objects)
 - Normal, star-forming disks
 - Low mass, LSB dwarfs
 - Potential for future star formation (HI content)
 - HI deficiency in clusters
 - History of tidal events
- **Rotational velocities**
 - Dark matter
 - Redshift-independent distances via Tully-Fisher relation

Sometimes...


- **HI absorption (not emission):**
 - "Optical depth"
 - Fundamental constant evolution



HI in M31




Credit: R. Braun



Hubble's Law

The dominant motion in the Universe is the smooth expansion known as the "Hubble flow".

Hubble's Law: $V_{\text{obs}} = H_0 D$
 where H_0 is Hubble's "constant" and D is distance in Mpc



Spread in velocity for objects in a cluster due to their orbital motion within the cluster.

$$\text{Recessional velocity} = \text{Hubble's constant} \times \text{Distance}$$

Relativistic Doppler Formula



- We observed galaxies/quasars with redshifts of $\sim 7-10$
- That does not mean that they are traveling faster than the speed of light

$$\frac{\text{shift in wavelength}}{\text{wavelength}} = \frac{\Delta\lambda}{\lambda} = \boxed{z} \quad \text{REDSHIFT}$$
$$z = \sqrt{\frac{\left(1 + \frac{v}{c}\right)}{\left(1 - \frac{v}{c}\right)}} - 1$$

This reduces to the simple Doppler formula for $v \ll c$.

For $z = 10$, this becomes

$$v = c \sqrt{1 - \left(\frac{1}{11}\right)^2} = 0.995 c$$

$$1 + z = \frac{1}{\sqrt{1 - v^2/c^2}}$$

$$v = c \sqrt{1 - \frac{1}{(1+z)^2}}$$

In fact, the Cosmic Microwave Background photons have a redshift $z = 1000$!

HI: The fuel for star formation




M81 – Spiral Galaxy (Type Sb)


Distance: 12,000,000 light-years (3.7 Mpc)

Image Size = 14 x 14 arcmin


Visual Magnitude = 6.9



X-Ray: ROSAT




Ultraviolet: ASTRO-1




Visible: DSS




Visible: R. Gendler




Near-Infrared: Spitzer



Mid-Infrared: Spitzer



Far-Infrared: Spitzer



HI
21 cm
line


http://coolcosmos.ipac.caltech.edu/cosmic_classroom/multiwavelength_astronomy/multiwavelength_museum/m81.html

9


UAT 17.06

ALFALFA





HI in NGC 2403



HI distribution



starlight

HI traces beyond the stellar disk => dark matter halo

HI traces interactions!




Gravitational interactions perturb preferentially the outer regions of the galaxies.


HI shows both the HI distribution and the velocity field.

HI is an excellent tracer of tidal interactions.


M81/M8/NGC3077






HI traces interactions!





M81/M8/NGC3077



HI traces interactions!



HI traces interactions!



15

UAT 17.06


ALFALFA



And sometimes HI is lost!



Virgo, A Laboratory for Studying Galaxy Evolution




The HI gas in galaxies moving through the hot intracluster medium of clusters of galaxies is stripped by the pressure of the hot gas


⇒ Ram pressure stripping

- Galaxies are HI deficient (lower than expected HI masses)
- HI disks of deficient galaxies are smaller in size (shrunk)

The ALFALFA population



Red: SDSS
Blue: ALFALFA



- 6500 sqd of high galactic latitude sky; median $cz \sim 8800$ km/s
- Undersamples clusters but traces well the lower density regions
- Large overlapping areas with SDSS and GALEX
- HI surveys detect the star-forming galaxy population
 - Nearly all star-forming galaxies contain HI

17





UAT 17.06

ALFALFA

The ALFALFA population

- Star-forming galaxies but not the red sequence



Shan Huang (Cornell) PhD thesis
Huang+(2012b) ApJ 756, 113

18

UAT 17.06

ALFALFA




HI-stellar scaling relations

Virtually all SF galaxies contain HI but the red sequence galaxies contain (none).

- **HI fraction falls** as the stellar luminosity/mass increases
- **Low luminosity** SF galaxies are HI-dominated (more mass in HI than in stars)
- **Increased scatter and break in slope** of M_{HI} versus M_{stellar} relation (and also f_{HI}) below $M_{\text{stellar}} \sim 10^9 M_{\odot}$
- HI represents the fuel **reservoir** for future star formation.

Huang+(2012b) ApJ 756, 113



HI cosmology





- The HI Mass Function: # galaxies per interval of HI mass per unit volume (analogous to a luminosity function) (Martin+2010 ApJ 723, 1359)
 - Cosmic density of HI at redshift 0
- The HI correlation function: how do HI galaxies cluster? (Martin+2012 ApJ 750, 38)
 - HI galaxies are the least-clustered population
- The HI velocity width function gives a perspective on the dark matter halo mass function. (Papastergis+ 2011 ApJ 739, 38)
 - We don't understand how gas/stars fit into halos
 - But gas richness is related to halo angular momentum

All of these yield insight into the distribution of dark matter halos, in this case ones which are gas-bearing, regardless of their stellar content.



HI-selected ALFALFA population

The HI population is **much less clustered** on small scales, but follows the DM on large scales.



The HI population is **least clustered known** => environmentally-driven processes are minimized.

Martin + (2012) Ap J 750, 38

A link between spin and gas richness





Halo spin parameter:

$$\lambda = J|E|^{1/2} G^{-1} M^{-5/2}$$

Observationally (not so easy) $\lambda = 21.8 \frac{R_d \text{ (kpc)}}{(V_{\text{rot}} \text{ [km/s]})^{3/2}}$

Higher gas fraction \Leftrightarrow high spin parameter of DM halo!



Shan Huang+ (2012b) ApJ 756, 113

22




UAT 17.06

ALFALFA

Low mass galaxies: Insights from ALFALFA



- At low M_{star} , the **baryonic gas fraction** $M_{\text{HI}} / (M_{\text{HI}} + M_{\text{star}})$ **approaches 1**.
- Many low HI mass dwarfs are LSB and patchy, so their stellar masses are uncertain.
- For low mass star forming galaxies, most of the baryonic mass is in HI!!!!



ALFALFA: Are there "dark galaxies"?



- In agreement with previous results, ALFALFA finds that **fewer than 2%** of (clearly extragalactic; not ALFALFA UCHVCs) HI sources cannot be identified with an optical counterpart.
- The majority of objects without OC's are **found near to** galaxies with similar redshifts.

Dark galaxies:

The burden is always on us to prove that

- (1) the signal is real and
- (2) there is no OC even at low surface brightness
- (3) the HI is not tidal in origin
- (4) not an OHM at $z \sim 0.2$

Luke Leisman, PhD thesis (Cornell)

Karen Lee-Waddell, PhD thesis (Queen's)

Steven Janowiecki, Bill Janesh PhD thesis (Indiana)

+ Cannon, Salzer, Rhode, Jozsa, Adams, Darling, RG, MH

ALFALFA
“(Almost) Dark”
galaxies project

24

UAT 17.06

ALFALFA



Why so much gas, so few stars?



HI1232+20 (a.k.a. "Coma P")


- Peak SB(g) 26.5 mag/sq"
- Visible in GALEX, not SDSS
- $\log M_{\text{HI}}/M_{\odot} = 8.83$ @25 Mpc
- $M_g \sim -12.7$ @25 Mpc
- $M_{\text{HI}}/L_g \sim 44$
- $R_{\text{HI}} \sim 23 \times 11$ kpc @ 25 Mpc
- $R_{\text{opt}} \sim 7.3 \times 3.7$ kpc @ 25 Mpc
- Hint of ordered rotation
- "Almost dark"; no H α

Does a different SFL hold?

Steven Janowiecki (IU)

Luke Leisman (Cornell)

+ Adams, Josza, Cannon, Salzer, Rhode, RG, MH



Janowiecki + 2015 ApJ 801, 96

Catie will tell us more! 25

UAT 17.06

ALFALFA



The $z \sim 0$ HI population from ALFALFA



- HI blind surveys do not “see”:
 - the “red sequence”
 - clusters (HI deficiency)
- HI-selected galaxies are the **least clustered** population (Martin+ 2012, Papastergis+2013)
- The ALFALFA population is typically **bluer**, of **lower metallicity** and lower **extinction**, consistent with having **extended disks** and lower SFEs.
- Galaxies with **higher HI fraction** are hosted in halos with **higher spin λ** (HI tells us about dark matter?????!!!)
- HI **dominates** the (visible) baryons in low mass galaxies.
- Some of the dwarfs are nearly dormant : could there be a population of low mass dwarfs with **stellar/gas contents so low** that we don't see them except when they accrete a small amount of gas and form a few stars (Leo P) =>
 - “Too Shy to Shine” = “Vanishing Cheshire Cat” (Salpeter & Hoffman 1995; Kormendy & Freeman 2015, astro-ph/1411.2170)

26



UAT 17.06

ALFALFA